

# Construction of a Novel DNA Vaccine Candidate encoding LmSTII1-PpSP42 Fusion Protein from *Leishmania major* and *Phlebotomus papatasi* Against Cutaneous Leishmaniasis

Touraj Miandoabi<sup>1</sup>, Fariborz Bahrami<sup>2</sup>, Vahideh Moein Vaziri<sup>1</sup>, Soheila Ajdary\*<sup>2</sup>

## Abstract

**Background:** Cutaneous leishmaniasis (CL) is a serious public health problem in many tropical countries. The infection is caused by a protozoan parasite of *Leishmania* genus transmitted by Phlebotominae sandflies. In the present study, we constructed a eukaryotic expression vector to produce a fusion protein containing LmSTII1 from *Leishmania major* (*L. major*) and PpSP42 from *Phlebotomus papatasi* (*Ph. papatasi*). In future studies we will test this construct as a DNA vaccine against zoonotic CL.

**Methods:** The nucleotide sequences encoding the LmSTII1 protein and a fragment encoding 79% of PpSP42 were amplified using *L. major* and *Ph. papatasi* genomic DNA, respectively. The amplicons were cloned into the pcDNA3.1(+) eukaryotic expression vector. The recombinant plasmid pcDNA-LmSTII1Pp42 was propagated in *Escherichia coli* (*E. coli*) and used to transfect HEK-293T cells. The expressed fusion protein was analyzed by Western blotting using anti-LmSTII1 mouse serum.

**Results:** Sequences encoding LmSTII1 and partial PpSP42 were cloned into pcDNA3.1(+). Production of the recombinant LmSTII1Pp42 fusion protein was confirmed by Western blotting.

**Conclusions:** An LmSTII1Pp42 fusion protein was expressed in HEK-293T cells. This construct may be an effective DNA vaccine against CL.

**Keywords:** Cloning, DNA vaccine, *Leishmania major*, LmSTII1, PpSP42.

## Introduction

Leishmaniasis are a group of important worldwide parasitic infectious diseases caused by obligatory intracellular protozoans of the genus *Leishmania*. These are transmitted to their vertebrate hosts, including humans, by the bite of infected female sandflies during blood feedings (1-3). Infections caused by *Leishmania* parasites manifest in a wide range of clinical forms in humans (4), including asymptomatic, cutaneous, diffuse cutaneous, mucosal cutaneous, visceral, and post-kala-azar dermal leishmaniasis (5). Cutaneous leishmaniasis (CL) is the most common form of the disease, caused by propagation of the parasite inside

the host's dermal macrophages. Two-thirds of all new CL cases occur in Afghanistan, Algeria, Brazil, Colombia, Iran, and Syria (1). Cutaneous leishmaniasis is highly prevalent in Iran; in 2012–2013, 37,001 cases were reported with the incidence estimated at 25/100,000 people (6).

Despite efforts by researchers worldwide, CL remains an important public health problem in many parts of Asia, North Africa, and Central and South Americas. Chemotherapeutics against the disease are currently available; however, due to toxicity, side effects, and high costs, they cannot be readily used. In addition, no effective vaccine has yet been

1: Department of Parasitology and Mycology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2: Pasteur Institute of Iran, Department of Immunology, 69 Pasteur Ave., Tehran, Iran.

\*Corresponding authors: Soheila Ajdary; Tel: +91 21 64112840; Fax: +91 21 64112840; E-mail: sohary@yahoo.com.

Received: Dec 12, 2017; Accepted: Jan 15, 2018

authorized against CL (7-9). In recent years, the complex relationship between the vector, the parasite, and the vertebrate host has been recognized. It has been shown that components in the vector's saliva contribute to the infection (10-12).

The induction of cellular immune responses, and more specifically a CD4<sup>+</sup> Th1 response, is essential to control the infection. A proper response against CL involves activation of the infected macrophages by secretion of Th1 cytokines including IFN- $\gamma$ . The parasites inside the phagolysosomes of the activated macrophages are then destroyed by cytotoxic reagents including hydrogen peroxide and nitric oxide (13). A protective CL vaccine should induce sufficient cellular immunity and cell responses to overcome the infection (14).

In lieu of safe, efficient, and affordable anti-leishmanial vaccines or drugs, new modes of prophylactics and therapies are essential to combat CL (9). In this regard, DNA vaccines have been shown to enhance long-lived humoral and cellular immune responses *in vivo* in a variety of animal models (15). Moreover, vaccination with plasmid DNA has been shown to induce protective immunity through both MHC class I- and class II-restricted T cell responses in a variety of experimental infection models, while in general, conventional protein vaccinations have been shown to induce only MHC class II-restricted responses (16). A potent immune response against CL is unlikely to be induced by only one antigen; therefore, different antigenic compounds need to be studied (17). Incorporation of such candidate antigens in plasmids encoding more than one antigen may enhance DNA vaccine immunogenicity (18). As the first step of an ongoing project with the aim of developing a DNA vaccine against CL, a recombinant plasmid with one antigen from the parasite and another from its vector was constructed. From *Leishmania major* (*L. major*), we used LmSTII, a homologue of yeast stress-inducible protein STII, which contains six copies of tetratricopeptide (TPR) consensus motif and is related to a highly-conserved family of stress-inducible proteins (19). Webb and colleagues in 1996 identified the full-length *L. major* LmSTII gene, which is expressed in a

temperature-dependent manner in both *L. major* promastigotes and amastigotes by a 26-37 °C shift in the culture temperature. Vaccination experiments with recombinant LmSTII protein plus either IL-12 or LmSTII DNA resulted in a mixed cellular response that was skewed toward a Th1 phenotype and protected BALB/c mice (19-22). We also used exon 2 of a gene that encodes 79% of *Ph. papatasi* salivary gland protein 42 (PpSP42) in the construct.

The sandfly's saliva contains proteins that modulate the mammalian host's immunological and physiological responses to the bites to accommodate parasitic invasion and establishment of the infection (23, 24). Different sandfly salivas differ in composition (25), and the induced immune responses are species-specific (26). PpSP42 is a homolog of LJM11 salivary gland protein of *Lutzomyia* (*Lu.*) *longipalpis*, the *L. brasiliensis* vector in the New World which its long-term protection against CL has been recently shown (27). Among the interesting attributes of this protein are its lack of homology with mammalian proteins, its capability to induce antibodies in humans and dogs, and its ability to be over-produced in prokaryotic expression systems (28, 29). PpSP42 was identified by Valenzuela and colleagues in 2001; however, its function remains to be characterized (23). Here, following construction of a plasmid to express LmSTIIp42, its production in mammalian cells was demonstrated using human embryonic kidney 293 (HEK) cells.

## Materials and methods

### Parasite

*L. major* promastigotes MRHO/IR/75/ER were cultured in RPMI 1640 medium (Biosera, France) supplemented with 10% heat-inactivated fetal calf serum (FCS; Biosera, France), 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin, and incubated at 24 °C. The promastigotes were harvested in mid-logarithmic phase at a density of  $2 \times 10^6$ /ml.

### Bioinformatics analyses

Nucleotide sequences were verified by Chromas Lite 2.01 (Technelysium Pty Ltd., Australia) and Sequencher 4.8 (Gene Codes, USA). The nucleotide-to-amino acid translations and the theoretical protein molecular weight estimations

were performed by ExPASy Bioinformatics Resource Portal <http://web.expasy.org/>.

### Genomic DNA extraction

Genomic *L. major* DNA was extracted from the promastigotes by phenol/chloroform method. Briefly,  $2 \times 10^7$  promastigotes were pelleted by centrifugation for 10 min at 800 x g, washed with phosphate buffered saline (PBS), and lysed in 350  $\mu$ l lysis buffer containing 0.1 M Tris-HCl, pH 8.0, 1% sodium dodecyl sulfate (SDS), 0.1 M NaCl, 10 mM EDTA, and 3.5  $\mu$ l of proteinase K (100  $\mu$ g/ml) at 55 °C for 2 h. The lysate was added to an equal volume of phenol/chloroform (450  $\mu$ l) to remove proteins. This mixture was centrifuged at 13,400 x g at 4 °C for 15 min and an equal volume of chloroform was added to the supernatant, which was then re-centrifuged as above. The supernatant was mixed with 1/10 volume of 3 M sodium acetate and two volumes of 100% ethanol to precipitate the DNA, and centrifuged as above for

10 min. The DNA pellet was washed with 70% ethanol, dissolved in 100  $\mu$ l of sterile distilled water, and stored at -20 °C until use. Genomic DNA from a female *Ph. papatasi* sandfly trapped in Kaleibar region of East Azerbaijan (Iran) and isolated by Dr. Parviz Parvizi, Department of Parasitology, Pasteur Institute of Iran, was obtained as a gift. The DNA concentrations and their quality were assessed by spectrophotometry on a NanoDrop 1000a (Thermo Scientific, USA) and electrophoresis on 1% agarose gels.

### PCR amplifications and sub-cloning of LmSTII and PpSP42


Promastigote genomic DNA was used as the template to PCR amplify the 1641-bp full-length LmSTII gene encoding a 62.1 kDa protein. Forward and reverse primers were designed based on DNA sequences of LmSTII available in GenBank by accession number U73845 (Table 1).

**Table 1.** PCR primers. The underlined segments depict the incorporated restriction sites. The bold letters on LmSTII F and PpSP42 R show a Kozak and a stop codon, respectively.

| Gene     | Primer (5' - 3')                                   |
|----------|--|
| LmSTII F | ATCTGGT <u>ACCCACCATGGCG</u> GACGCAACTGAGCTGAAGAAC |
| LmSTII R | GGTCGAATTCCTGACCAAAACGAATGATGCCAGCTG               |
| PpSP42 F | GCTCGAATTCGCTTACGATTCAGGAAATATTGTACCAAC            |
| PpSP42 R | ATCACTCGAGTCACATAATGTCTGTGCCAAAATTGAAG             |

The forward primer contained a *Kpn*I (GGTACC) restriction site and a Kozak translation initiation sequence consensus (CACCATGGCG). The reverse primer contained an *Eco*RI (GAATTC) restriction site. To amplify PpSP42, *Ph. papatasi* genomic DNA was used as the template along with

primers, designed using DNA sequences available in GenBank by accession number KX611849.1 (Table 1). The lack of introns between the primers and the exons was verified by comparing *Ph. papatasi* sp42 mRNA (GenBank accession number AF335491.1) with the *Ph. Papatasi* genome sequence (NCBI PRJNA20293; Fig. 1).



**Fig. 1.** Schematic view of *Ph. Papatasi* PpSP42 gene (GenBank accession number AF335491) exons and introns, depicting the locations of the primers on exon 2.

The forward and reverse primers, containing *EcoRI* (GAATTC) and *XhoI* (CTCGAG) restriction sites, respectively, and a stop codon (TCA) in the reverse primer, were used to amplify a 945-bp amplicon. The amplifications were performed in 25  $\mu$ l volumes containing 1  $\mu$ l of genomic DNA as the template, 10 pmol of each primer, 1.5 mM MgCl<sub>2</sub>, and ExPrime Taq™ DNA Polymerase (Genet Bio, Republic of Korea). The thermocycling program was 94 °C for 10 min initial denaturation, followed by 30 cycles of 94 °C for 1 min, 60 °C for 1 min, and 72 °C for 2 min, with a final extension at 72 °C for 10 min. The amplicons were electrophoresed on 1% agarose gels, stained with DNA Green viewer (Pars Tous, Iran), and visualized on a UV transilluminator. The size markers used to estimate PCR products were 100-bp and 1-kbp DNA ladders (SinaClon, Iran).

The LmSTII and Pp42 amplicons were purified with a PrimePrep™ PCR Purification Kit (GeNet Bio, South Korea) and ligated into the pTZ57R/T cloning vector (InsTAclone™ PCR product cloning kit, Thermo Scientific, USA) following the manufacturer's protocol. The resulting colonies were screened by colony-touch PCR using the above primer sets specific to each amplicon (Table 1). Extracted plasmids were prepared from the TA-cloned plasmids using a TIANprep kit (Tiangen, China). The integrities of pT-LmSTII and pT-PpSP42 plasmids were verified by nucleotide sequencing using M13F and M13R universal primers from Gen Fanavaran Co. (Iran).

#### **Construction of eukaryotic expression vector for the fusion protein**

LmSTII was cloned into pcNDA3.1 (+) eukaryotic expression vector to produce pCLmSTII as follows: the TA-cloned pT-LmSTII plasmid and pcDNA3.1 (+) were double-digested separately with *KpnI* and *EcoRI* (Fermentas, Germany) in 1X Tango buffer. The double-digestion reactions were performed in two steps. First, in 40  $\mu$ l volumes, 200 ng of pT-LmSTII or pcDNA3.1 (+) were incubated with 4  $\mu$ l of *KpnI* at 37 °C for 2 h. Second, 1  $\mu$ l of *EcoRI* was added to each reaction and the mixtures were incubated at 37 °C for an additional 2 h. The digested products were analyzed by electrophoresis on 1% agarose gels as

described above. The bands belonging to the double-digested LmSTII and the vector fragments were extracted from the agarose gel with an AccuPrep® Gel Purification Kit (Bioneer, Germany). The fragments were then ligated in a 1:1 molar ratio using T4 DNA ligase (Vivantis, Malaysia). *Escherichia coli* Top10 cells were transformed with the preliminary recombinant plasmid, named pCLmSTII. Colonies grown on agar plates containing 50  $\mu$ g/ml of ampicillin were screened by colony-touch PCR using LmSTII-specific primers. The expected amplicon size was 1641 bp. A selected colony was grown overnight and pCLmSTII was extracted as described above. The plasmids were analyzed by *EcoRI* single- and *EcoRI/KpnI*-double-digestions. The expected linear plasmid sizes from the single digestion were 7068 bp, and for the *EcoRI/KpnI* double-digestion, 1640 bp for the LmSTII insert and 5428 bp for pcDNA3.1(+). The integrity of the construct and the lack of deletions or mutations were verified by nucleotide sequencing (Gen Fanavaran Co., Iran).

Next, PpSP42 was added to pCLmSTII to produce pCLmSTII-Pp42 as follows: the recombinant plasmids PpSP42 and pCLmSTII were double-digested at 37 °C for 2 h with 1  $\mu$ l each of *XhoI* and *EcoRI* (Fermentas, Germany) in separate 20  $\mu$ l volumes containing 2  $\mu$ g of DNA and 1X Tango buffer. The digestion products of expected fragment sizes of 939 bp for PpSP42 and 7064 bp for pCLmSTII were gel-extracted and ligated as above. The recombinant fusion plasmid, named pCLmSTII-Pp42, was used to transform competent *E. coli* Top10 cells. Colonies were selected as above and recombinants were identified by *EcoRI* digestion with expected linear plasmid size of 7950 bp and *EcoRI/XhoI* double digestion with expected fragment sizes of 2595 bp for the LmSTII-Pp42 insert and 5428 bp for pcDNA3.1(+), followed by DNA sequencing as described above.

#### **Transfection of HEK-293T eukaryotic cells with the recombinant plasmid**

HEK-293T cells were transfected with pCLmSTII-Pp42 using X-tremeGENE HP DNA Transfection Reagent (Roche, Germany). After two passages the cells were seeded at  $4 \times 10^5$

cells/2 ml medium/well in two wells of a six-well plate with 9.4 cm<sup>2</sup> diameter wells, and grown in RPMI 1640 medium (Biosera, France) supplemented with 10% FCS (Biosera, France), 1% glutamine, and 0.1% penicillin/streptomycin in aseptic conditions, and incubated at 37 °C in 5% CO<sub>2</sub> with 70% humidity for 48 h. The transfection reagents for each well contained 200 µl of serum-free RPMI medium, 6 µl of transfection reagent (1:3 ratio), and 4 µg of either pCLmSTIIPp42 or pcDNA3.1 (+) as a control. After mixing, the plate was incubated for 15 min at room temperature and the transfection complexes were added dropwise to the 60-70% confluent cells with gentle shaking to ensure even distribution of the ingredients in each well. The plate was incubated for 72 h and protein expression was assayed.

#### **Extraction of recombinant proteins, SDS-PAGE, and Western-blotting**

Following incubation with the transfection reagents, the cells were harvested and washed three times with sterile PBS and centrifugation at 13,400 x g at 4 °C for 1 min. The cells were then lysed with 300 µl of NP40 non-denaturing lysis buffer containing 150 mM NaCl, 50mM Tris-HCl, pH 8.0, 1% NP-40, and protease inhibitors, and incubated for 30 min at room temperature. The cell lysates were vigorously vortexed for 10 min and then centrifuged at 13,400 x g at 4 °C for 20 min. The clear cell lysate supernatants were collected and stored at -80 °C until use.

Proteins were separated by 8% SDS-PAGE using 100 µl of the clear lysates along with pre-stained protein ladder SM7002 (SinaClon, Iran)

at 40 V for 2 h. The gel was stained overnight with Coomassie Blue staining solution. An additional SDS-PAGE was performed and the polypeptides from the gel were transferred to a PVDF membrane (Millipore, USA) at 15 V for 30 min. The membrane was blocked with PBS containing 1% BSA (Sigma, Germany) for 2 h and then incubated overnight at 4 °C with a polyclonal murine anti-LmSTII antibody diluted 1/100. After washing, the membrane was incubated for 2 h with anti-mouse IgG-HRP (Sigma, Germany). The proteins were detected with DAB (Bio-Rad, Germany) staining.


## **Results**

### **Construction of the DNA vaccine**

A 1641-bp *L. major* LmSTII amplicon and a 930-bp *Ph. papatasi* PpSP42 amplicon were obtained by PCR (Fig. 2). The predicted sizes of the cloned fragments were verified by double-digestions of pCLmSTIIPp42 with *Kpn1/EcoR1* and *Kpn1/Xho1* (Fig. 3), and the integrities of pCLmSTII and pCLmSTIIPp42 were verified by DNA sequencing. Based on the sequencing results, the expressed pCLmSTIIPp42 fusion protein contains 859 amino acids (Fig. 4).


### **Expression and detection of LmSTIIPp42 fusion protein in eukaryotic cells**

HEK-293T cells were transfected with pCLmSTIIPp42 and expression of the fusion protein with an approximate Mw of 100 kDa (theoretical Mw: 97.1 kDa) was verified by SDS-PAGE and Western blotting with a polyclonal anti-LMSTII antibody (Fig. 5).



**Fig. 2.** A: Agarose gel electrophoresis of A: the full length *L. major* LmSTII gene fragment (expected size of amplicon: 1641bp), B: *Ph. papatasi* partial PpSP42 fragment (expected size of amplicon: 930 bp).






**Fig. 3.** A: Schematic diagram of pCLmSTI1Pp42 construct composed of pcDNA3.1(+) eukaryotic expression vector and in-frame *L. major* LMSTI1 and *Ph. Papatasi* SP42 genes. B: pCLmSTI1 double-digested with *KpnI* and *EcoRI* (expected digested fragments: ~5400 bp and ~1600 bp). C: pCLmSTI1Pp42 double-digested with *KpnI* and *XhoI* (expected digested fragments: ~5400 bp and ~2500 bp).

```

1  MADATELKNK  GNEEFSAGRY  VEAVNYFSKA  IQLDEQNSVL  YSNRSACFAA  MQKYKDALDD
61  ADKCISIKPN  WAKGYVRRGA  ALHGMRRYDD  AIAAYEKGLK  VDPSNSGCAQ  GVKDVQVAKA
121 REARDPIARV  FTPEAFRKIQ  ENPKLSLLML  QPDYVKMVDT  VIRDPSQGR L  YMEDQRFALT
181 LMYLSGMKIP  NDGDGEEEEER  PSAKAAETAK  PKEEKPLTDN  EKEALALKEE  GNKLYLSKKF
241 EEALTKYQEA  QVKDPNNTLY  ILNVSAVYFE  QGDYDKCIAE  CEHGIEHGRE  NHCDYTIIAK
301 LMTRNALCLQ  RQRKYEA AID  LYKRALVEWR  NPDTLK KLE  TE  CEKEHQKAVE  EAYIDPEIAK
361 QKKDEGNQYF  KEDKFPEAVA  AYTEAIKRN P  AEHTSYSNRA  AAYIKLGAFN  DALKDAEKCI
421 ELKPDFVKGY  ARKGHG YFWT  KQYNRALQAY  DEGLKVDPSN  ADCKDGRYRT  IMKIQEMASG
481 QSADGDEAAR  RAMDDPEIAA  IMQDSYMQLV  LKEMQNDPTR  IQEYMKDSGI  SSKINKLISA
541 GIIRFGQEF A  YDSGNI VPTG  VAYDAASKML  FFGIPRKYSK  VPITLAQLST  RSYNSAEIPN
601 PPLDKFSGKS  KQPLSSVYQP  VIDDCRRLWV  LDVGIVENEA  ERKTYPIIKP  SLIAFDLTKS
661 NYPEIHRYEL  TGEAGKNPLG  YGGFAADV N  PKRCS DKNEK  TYIYIANFDE  NSLIVYDKKK
721 GEAWSLKDDS  FKPEGVTTFT  LNGKEHKFKA  GIFGIALGDR  NKEGNRPAYY  LAGSSTKLYR
781 LDTKLLKKKS  SKLEPKLIGD  RGFKTEAIAL  AHPETKVL F  FAEADSRQVS  CWNIKHELKP
841 ENVGVIYTNP  NFNFGTDIM
    
```

**Fig. 4.** Predicted amino acid sequence of the pCLmSTI1Pp42 fusion protein. The bold residues represent LmSTI1 and the underlined residues represent the partial PpSP42 protein sequence. The highlighted grey residues are encoded by pcDNA3.1(+).



**Fig. 5.** A: SDS-PAGE of clear lysate of pCLmSTI1Pp42-transfected HEK-293T cells, after 72 h of incubation. B: Western blot of clear lysate of pCLmSTI1Pp42-transfected HEK-293T cells after 72 h of incubation. Protein was detected by polyclonal murine anti-LmSTI1 antibody

### Discussion

Leishmaniasis remains a major health problem in many countries worldwide and the infection rate has remained steady in recent decades. Although limiting the reservoirs and vectors as well as identifying new drugs offers promise, particularly in view of the emerging drug resistances, development of safe and effective vaccines remains the best hope for achieving a definitive control of leishmaniasis (30). The major steps to increase the efficiency of a new vaccine against CL are to identify suitable new antigens and adjuvants and recognize the type of immune response generated by the protected hosts (31). Achieving an effective vaccine against CL should therefore trigger Th1 immune responses, which could potentially be due

to recognition of specific antigens from both the vector and parasite.

One of the main strategies to control CL is to develop a DNA vaccine. The immune responses against DNA vaccines can induce both CD4+ and CD8+ T cells, which leads to polarized cellular immunity, manifested by activated Th1 cell signaling, similar to an adjuvant (32). In contrast, production of a Th2 immune response is associated with susceptibility to, and intensification of, the disease (33). In this regard, various antigens including TSA, KMPII, TRYP, LACK, GP63, KMP-11, CPA, CPB, PSA2, H1, Leish-111f, and P4 have been evaluated as parts of potential DNA or recombinant vaccines against leishmaniasis (22, 34-41). Among the antigens of interest, LmSTII, which is constitutively expressed in both *L. major* promastigotes and amastigotes, has been proposed as a vaccine candidate against cutaneous and visceral leishmaniasis (20, 22). Moreover, several *Ph. papatasi* and *Lu. longipalpis* salivary proteins have been characterized. Of these, SP15 and

LJM19 have been used in DNA vaccine assays with various degrees of success (23, 42-44).

Here, we expressed a recombinant eukaryotic plasmid containing the *L. major* LmSTII gene in conjunction with exon 2 of the *Ph. papatasi* PpSP42 gene in HEK-293T eukaryotic cells. The potential protection by this vaccine candidate against experimental *L. major* infection will be evaluated in our laboratory in an animal model.

### Acknowledgment

This project was funded by research grant # IPI-726 from the Pasteur Institute of Iran. The authors thank Dr. P. Parvizi, Department of Parasitology, Pasteur Institute of Iran, for providing *Ph. papatasi* genomic DNA. We also thank Dr. F Riazi-Rad, Department of Immunology, Pasteur Institute of Iran, for his technical assistance. This article has been extracted from the Ph.D. thesis written by T. Miandoabi (registration no. 219) from the Department of Medical Parasitology and Mycology, School of Medicine at Shahid Beheshti University of Medical Sciences.

### References

1. Bates PA. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies. *International journal for parasitology*. 2007; 37(10):1097-106.
2. Alvar J, Vélez ID, Bern C, et al. Leishmaniasis worldwide and global estimates of its incidence. *PloS one*. 2012;7(5):e35671.
3. Masina S, M MG, Demotz SO, et al. Protection against cutaneous leishmaniasis in outbred vervet monkeys using a recombinant histone H1 antigen. *The Journal of infectious diseases*. 2003;188(8):1250-7.
4. Badirzadeh A, Taheri T, Abedi-Astaneh F, et al. Arginase activity of *Leishmania* isolated from patients with cutaneous leishmaniasis. *Parasite immunology*. 2017; 39(9):e12454-n/a.
5. Badirzadeh A, Mohebbali M, Ghasemian M, et al. Cutaneous and post kala-azar dermal leishmaniasis caused by *Leishmania infantum* in endemic areas of visceral leishmaniasis, northwestern Iran 2002-2011: a case series. *Pathogens and global health*. 2013;107(4):194-7.
6. Badirzadeh A, Mohebbali M, Asadgol Z, et al. The burden of leishmaniasis in Iran, acquired from the global burden of disease during 1990-2010. *Asian Pacific Journal of Tropical Disease*. 2017;7(9):513-8.
7. Sundar S, More DK, Singh MK, et al. Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clinical infectious diseases*. 2000;31(4):1104-7.
8. Zarrati S, Mahdavi M, Tabatabaie F. Immune responses in DNA vaccine formulated with PMMA following immunization and after challenge with *Leishmania major*. *Journal of parasitic diseases*. 2016;40(2):427-35.
9. Montakhab-Yeganeh H, Abdossamadi Z, Zahedifard F, et al. *Leishmania tarentolae* expressing CXCL-10 as an efficient immunotherapy approach against *L. major*-infected BALB/c mice. *Parasite Immunology*. 2017; 39(10):e12461
10. Ribeiro J. Role of saliva in blood-feeding by arthropods. *Annual review of entomology*. 1987;32(1):463-78.

11. Titus RG, Ribeiro J. Salivary gland lysates from the sand fly *Lutzomyia longipalpis* enhance *Leishmania* infectivity. *Science*. 1988;239(4845):1306-8.
12. Ribeiro J. Blood-feeding arthropods: live syringes or invertebrate pharmacologists? *Infectious agents and disease*. 1995;4(3):143-52.
13. Pirdel L, Zavarán Hosseini A, Rasouli M. Immune response in susceptible BALB/c mice immunized with DNA encoding Lipophosphoglycan 3 of *Leishmania infantum*. *Parasite immunology*. 2014;36(12):700-7.
14. Chen X-T, Chen L-L, Tan J-Y, Shi D-H, et al. Th17 and Th1 lymphocytes are correlated with chronic periodontitis. *Immunological investigations*. 2016;45(3):243-54.
15. Robinson HL. DNA vaccines: basic mechanism and immune responses. *International journal of molecular medicine*. 1999;4(5):549-604.
16. Gurunathan S, Stobie L, Prussin C, et al. Requirements for the maintenance of Th1 immunity in vivo following DNA vaccination: a potential immunoregulatory role for CD8+ T cells. *Journal of immunology*. 2000;165(2):915-24.
17. Coler RN, Reed SG. Second-generation vaccines against leishmaniasis. *Trends in parasitology*. 2005;21(5):244-9.
18. Dumonteil E. DNA vaccines against protozoan parasites: advances and challenges. *BioMed research international*. 2007.
19. Webb JR, Kaufmann D, Campos-Neto A, et al. Molecular cloning of a novel protein antigen of *Leishmania major* that elicits a potent immune response in experimental murine leishmaniasis. *The Journal of Immunology*. 1996;157(11):5034-41.
20. Webb JR, Campos-Neto A, Skeiky YA, et al. Molecular characterization of the heat-inducible LmSTII protein of *Leishmania major*. *Molecular and biochemical parasitology*. 1997;89(2):179-93.
21. Campos-Neto A, Porrozzi R, Greeson K, et al. Protection against cutaneous leishmaniasis induced by recombinant antigens in murine and nonhuman primate models of the human disease. *Infection and immunity*. 2001;69(6):4103-8.
22. Campos-Neto A, Webb JR, Greeson K, et al. Vaccination with plasmid DNA encoding TSA/LmSTII leishmanial fusion proteins confers protection against *Leishmania major* infection in susceptible BALB/c mice. *Infection and immunity*. 2002;70(6):2828-36.
23. Valenzuela JG, Belkaid Y, Garfield MK, et al. Toward a defined anti-*Leishmania* vaccine targeting vector antigens: characterization of a protective salivary protein. *The Journal of experimental medicine*. 2001;194(3):331-42.
24. Oliveira F, Jochim RC, Valenzuela JG, et al. Sand flies, *Leishmania*, and transcriptome-borne solutions. *Parasitology international*. 2009;58(1):1-5.
25. Ribeiro JM, Mans BJ, Arcà B. An insight into the sialome of blood-feeding Nematocera. *Insect biochemistry and molecular biology*. 2010;40(11):767-84.
26. Thiakaki M, Rohousova I, Volfova V, et al. Sand fly specificity of saliva-mediated protective immunity in *Leishmania amazonensis*-BALB/c mouse model. *Microbes and infection*. 2005;7(4):760-6.
27. Gomes R, Oliveira F, Teixeira C, et al. Immunity to sand fly salivary protein LJM11 modulates host response to vector-transmitted leishmania conferring ulcer-free protection. *J Invest Dermatol*. 2012;132(12):2735-43.
28. Teixeira C, Gomes R, Collin N, et al. Discovery of markers of exposure specific to bites of *Lutzomyia longipalpis*, the vector of *Leishmania infantum chagasi* in Latin America. *PLoS Negl Trop Dis*. 2010;4(3):e638.
29. Xu X, Oliveira F, Chang BW, et al. Structure and function of a "yellow" protein from saliva of the sand fly *Lutzomyia longipalpis* that confers protective immunity against *Leishmania major* infection. *J Biol Chem*. 2011;286(37):32383-93.
30. Mauel J. Vaccination against *Leishmania* infections. *Current Drug Targets-Immune, Endocrine & Metabolic Disorders*. 2002;2(3):201-26.



31. Von Stebut E, Udey MC. Requirements for Th1-dependent immunity against infection with *Leishmania major*. *Microbes and infection*. 2004;6(12):1102-9.
32. Ertl PF, Thomsen LL. Technical issues in construction of nucleic acid vaccines. *Methods*. 2003;31(3):199-206.
33. Melby PC. Vaccination against cutaneous leishmaniasis. *American journal of clinical dermatology*. 2002;3(8):557-70.
34. Skeiky YA, Coler RN, Brannon M, et al. Protective efficacy of a tandemly linked, multi-subunit recombinant leishmanial vaccine (Leish-111f) formulated in MPL adjuvant. *Vaccine*. 2002;20(27-28):3292-303.
35. Shaddel M, Oormazdi H, Akhlaghi L, et al. Cloning of *Leishmania major* P4 gene. *Tropical Medicine & International Health*. 2007;12:213.
36. Rodriguez-Cortes A, Ojeda A, Lopez-Fuertes L, et al. Vaccination with plasmid DNA encoding KMP11, TRYP, LACK and GP63 does not protect dogs against *Leishmania infantum* experimental challenge. *Vaccine*. 2007;25(46):7962-71.
37. Rafati S, Kariminia A, Seyde-Eslami S, et al. Recombinant cysteine proteinases-based vaccines against *Leishmania major* in BALB/c mice: the partial protection relies on interferon gamma producing CD8(+) T lymphocyte activation. *Vaccine*. 2002;20(19-20):2439-47.
38. Iborra S, Soto M, Carrion J, Alonso C, et al. Vaccination with a plasmid DNA cocktail encoding the nucleosomal histones of *Leishmania* confers protection against murine cutaneous leishmaniasis. *Vaccine*. 2004;22(29-30):3865-76.
39. Dondji B, Pérez-Jimenez E, Goldsmith-Pestana K, et al. Heterologous prime-boost vaccination with the LACK antigen protects against murine visceral leishmaniasis. *Infection and immunity*. 2005;73(8):5286-9.
40. Bhaumik S, Basu R, Sen S, et al. KMP-11 DNA immunization significantly protects against *L. donovani* infection but requires exogenous IL-12 as an adjuvant for comparable protection against *L. major*. *Vaccine*. 2009;27(9):1306-16.
41. Tabatabaie F, Dalimi A, Sharifi Z, et al. Cloning and sequencing of *Leishmania major* thiol-specific-antioxidant antigen (TSA) gene. *Iranian journal of parasitology*. 2007;2(4):30-41.
42. Zahedifard F, Gholami E, Taheri T, et al. Enhanced protective efficacy of nonpathogenic recombinant *Leishmania tarentolae* expressing cysteine proteinases combined with a sand fly salivary antigen. *PLoS neglected tropical diseases*. 2014;8(3):e2751.
43. Katebi A, Gholami E, Taheri T, et al. *Leishmania tarentolae* secreting the sand fly salivary antigen PpSP15 confers protection against *Leishmania major* infection in a susceptible BALB/c mice model. *Molecular immunology*. 2015;67(2 Pt B):501-11.
44. Gomes R, Teixeira C, Teixeira MJ, et al. Immunity to a salivary protein of a sand fly vector protects against the fatal outcome of visceral leishmaniasis in a hamster model. *Proceedings of the National Academy of Sciences*. 2008;105(22):7845-50.