

Inhibitory and Apoptotic Effects of Mannan-Mitomycin C Conjugate Against Transitional Cell Carcinoma and Normal Mouse Fibroblasts

Ladan Kabiri¹, Shiva Irani¹, Amin Reza Nikpoor², Jalil Tavakkol Afshari^{*2}

Abstract

Background: Many studies have shown the anticancer effects of mannan and mitomycin C on tumor cells. In this regard, the aim of this study was to investigate the inhibitory and apoptotic effects of a mannan-mitomycin-C conjugate on transitional cell carcinoma (TCC) and normal mouse L929 fibroblast cells.

Methods: The conjugate was synthesized according to previous studies. Both cell lines were cultured and the cytotoxic and apoptotic effects of the compounds in different concentrations were assessed using MTT and flow cytometry, respectively. The mannan-mitomycin C conjugate inhibited proliferation of both cell lines in time and concentration -dependent manners.

Results: The conjugate inhibited TCC cell proliferation more than that of L929 cells. Mitomycin C alone inhibited proliferation of both cell lines in both time and concentration -dependent manners, and the effect was greater on L929 than on TCC cells. Mannan had a relatively low inhibitory effect on TCC and no significant effect on L929 cells. The percentage of apoptosis was greater in TCCs than in L929 cells at the highest concentration of conjugate. Mitomycin C induced apoptosis more extensively in L929 cells than in TCC cells at 25 and 400 µg/ml. The effect of mannan was similar on both cell lines.

Conclusions: The mannan-mitomycin C conjugate has greater inhibitory and apoptotic effects on TCC than on L929 cells and may inhibit TCC.

Keywords: Apoptosis, Inhibitory Effect, Mannan-Mitomycin C Conjugate, MTT, TCC.

Introduction

One of the most common genitourinary malignancies is bladder cancer (1). More than 90% of bladder malignancies are associated with transitional cell carcinoma (TCC) (2). Several factors can cause malignant cells in the bladder, including smoking, occupational exposure, chronic inflammation in the bladder due to schistosomiasis, or factors such as age, gender, race, and genetics (3, 4). Methods used to treat bladder cancer include transurethral resection of bladder (TURB), chemotherapy, Bacillus Calmette-Guérin (BCG) therapy, and cystectomy (5, 6). Chemotherapy with thiotepa, doxorubicin, and mitomycin is common. Although these methods lead to a relative improvement in cancer patients' conditions, the

percentage of definitive treatment of the disease is low. Therefore, the use of complementary therapies, such as immunotherapy can help the treatment process (7). The most common bladder cancer immunotherapy treatment is intracellular BCG. One component of BCG is mannan (8). BCG induces immunological changes in bladder cells and stimulates the production of chemokines such as IL-8 and inflammatory cytokines (7).

Mannan is a polysaccharide found in the outer layers of yeast cell walls (9). The composition of mannan extracted from yeast *Saccharomyces (S.) cerevisiae* can be specifically linked to a receptor called toll-like receptor 4 (TLR4) from the pattern recognition

1: Department of biology, Science and Research Branch, Islamic Azad University, Tehran, Iran.

2: Department of Immunogenetic, Immunology Research Center, Bu-Ali Research Institute, Mashhad University of Medical Sciences

*Corresponding author: Jalil Tavakkol Afshari; Tel: +98 51 37112471, Fax: +98 51 37112596, E-mail: tavakolaj@mums.ac.ir.

Received: 10 Jan, 2017; Accepted: Feb 7, 2018

receptors family (10, 11) which were originally found on immune cell membranes (10, 12) and also exist on bladder epithelial cells (12, 13). The binding of mannan with this receptor induces the production of inflammatory cytokines such as interleukins (ILs) 6, 8, and 12 and tumor necrosis factor- α (TNF α), and reduces or increases tumor growth (14, 15). Mannan also contributes to immune function by affecting monocytes and stimulating the production of interleukins and TNF α (11).

Mitomycin C, an antibiotic derived from *Streptomyces Caspitosis* (16), is used to treat many cancers, including bladder cancer. Unfortunately its side effects include anorexia, fatigue, hemolytic-uremic syndrome, mucositis, myelosuppression, thrombocytopenia, and renal failure (17).

In this study, we investigated the *in vitro* effects of mannan-mitomycin C conjugate, and mannan and mitomycin C alone on the proliferation and apoptosis of TCC and L929 cells, which may help us to develop new drugs that are more effective against cancer cells than normal cells and have fewer side effects than currently available drugs.

Materials and methods

Cell culture

The TCC and L929 cell lines were purchased from Ferdowsi University Cell Bank (Mashhad). The cells were cultured at 37°C in a humidified 5% CO₂ atmosphere in Dulbecco's Modified Eagle Medium (DMEM)-high glucose containing 10% fetal bovine serum (FBS) (Gibco) and 1% penicillin/streptomycin.

Preparation and analysis of mannan-mitomycin C conjugate

The mannan-mitomycin C conjugate was synthesized according to the modified Matsumoto's method (18). In the first step, mannan from *S. cerevisiae* wild type strain was activated with cyanogen bromide. For this purpose, 0.1 g of mannan was dissolved in 10 mL of water. 55 mg of cyanogen bromide was added and the pH of the solution was adjusted to 10.7 with 1 M NaOH. Then, 0.1 g of 6-aminohexanoic acid was added and the pH was adjusted to 9 with 1 M HCl. To make the coupling reaction, the

solution was stirred at room temperature for 24 hr. In the next step, the mannan-6-aminohexanoic acid product was dialyzed against sodium carbonate (Na₂CO₃) at pH 9 in dialysis tubes with 12000-MW cut-off for 24 hr. Then, 10 mg of mitomycin C was dissolved in mannan-6-aminohexanoic acid solution and 0.2 g of 1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide hydrochloride was added. The pH was adjusted to 5.0-6.0 and the solution was stirred at room temperature for 24 hr. Finally, the solution was dialyzed for 24 hr. The amount of mitomycin C bound in the mannan-mitomycin C conjugate was determined by measuring the absorbance of mitomycin C at 364 nm on a spectrophotometer (Analytikjena, Germany).

MTT assay

The inhibitory effect of the mannan-mitomycin C conjugate and mannan and mitomycin C alone on TCC and L929 cell lines was measured by MTT assay. For this purpose, 20,000 cells per well were cultured in 96-well plates. Cells were incubated for 24 hr to allow them to attach to the plate. After this time, the TCC and L929 cells were incubated with 700, 350, 175, 87.5, 43.75, or 0 μ g/mL mannan-mitomycin C conjugate, 400, 200, 100, 50, 25, or 0 μ g/mL mitomycin C, and 10000, 5000, 2500, 1250, 625, or 0 μ g/mL mannan for 24, 48, or 72 hr. At the end of the incubations the supernatants were removed and 20 μ L of MTT solution was added. After 4 hr of incubation at 37 °C, 100 μ L of DMSO and 10 μ L of glycine buffer was added to each well. Absorbance of each well was measured on an ELISA reader (BioTek, USA) at a wavelength of 570 nm. All experiments were performed in triplicate. The results were reported as survival percentages at the various sample concentrations.

Cell Apoptosis Analysis

To determine cell apoptosis after treatment with mannan, mitomycin C, and the mannan-mitomycin C conjugate, flow cytometry with Propidium iodide (PI) and Annexin-V FITC was performed. For this purpose, 2×10^5 L929 and TCC cells per well were cultured in 6-well plates. After 24 hr of incubation to allow the cells to adhere to the wells, they were treated for 72 hr with 2 mL of the

conjugate at 700 µg/ml, 2 mL of mitomycin C at 400 µg/ml, 2 mL of mitomycin C at 25 µg/ml, 2 mL of mannan at 1250 µg/ml, and a well for each cell line with no treatment as a negative control. The cells were then washed with PBS and trypsinized. The cell suspension was centrifuged and the cell pellet was washed 2x with PBS containing 1% FBS. The cells were then solubilized in 100 µl of ready-to-use binding buffer (cat # 422201). In the next step, 5 µl of Annexin-V FITC and 10 µl of PI solution were added to each tube. The solubilized cells were gently vortexed and incubated for 15 minutes at room temperature in darkness. Finally, 400 µl of binding buffer was added to each tube and apoptosis was analyzed on a BD FACSCalibur (USA).

Morphological studies

After treatment of the L929 and TCC cells with the mannan-mitomycin C conjugate, mitomycin C, or mannan at different concentrations for 24, 48, and 72 hr, the cell morphologies were examined by inverse optical microscopy.

Statistical analysis

The results were analyzed using GraphPad Prism 6 and one-way analysis of variance (ANOVA) and Tukey's post-HOC test. In all tests, $P < 0.05$ was considered statistically significant.

Results

MTT results 24, 48, and 72 hr after adding mannan-mitomycin C conjugate to L929 and TCC cells

After 24 hr, no significant difference in cell viability was seen between the L929 and TCC cells treated either without or with the conjugate. The conjugate caused a slight decrease in viability of both the TCC and L929 cells, although this decrease was not significant ($P > 0.05$) (Fig. 1). After 48 hr of incubation with the conjugate however, viability was significantly less in the TCC than in the L929 cells ($P < 0.0001$). Cell viability decreased with increasing conjugate concentration. The mean (CI 95%) IC₅₀ for the TCC cells was 457.9 (262.3 to 799.3) µg/ml, while the effect of conjugate was not sufficient to kill 50% of the L929 cells. Cell viability was significantly less in the TCC cells than in the L929 cells at all conjugate concentrations ($P < 0.0001$). After 72 hr cell survival decreased for both cell lines with increasing conjugate concentrations and was significantly less in the TCC than in the L929 cells at conjugate concentrations of 175, 350, and 700 µg/mL ($P < 0.05$, 0.001, and 0.0001, respectively). The mean (CI 95%) IC₅₀ for the TCC cells was 187.0 (173.3 to 201.0) µg/ml, while in the L929 cell line, the mean (CI 95%) IC₅₀ was 551.5 (427.2 to 712.0) µg/ml.




Fig. 1. Growth inhibitory effect of mannan-mitomycin C conjugate on L929 and TCC cells. Cells were grown in 96-well plates and incubated for 24, 48, or 72 hr with 0, 43.75, 87.5, 175, 350 or 700 µg/mL of the mannan-mitomycin conjugate. Cell viability was measured with MTT assays. Mean $\pm$ SD, ANOVA, $n=3$. ($P < 0.05^*$, $P < 0.01^{**}$, $P < 0.001^{***}$, $P < 0.0001^{****}$).

MTT results 24, 48, and 72 hr after adding mitomycin C to L929 and TCC cells

After 24 hr of mitomycin C treatment, viability decreased significantly in both cell lines ($P < 0.0001$). The mean (CI 95%) IC₅₀s were 74.8 (63.8 to 87.6) µg/mL and 57.3 (38.6 to 85.1)

µg/mL for the TCC and L929 cells, respectively. After 48 hr the decreasing trend in survival rate was statistically significant ($P < 0.0001$) (Fig. 2, 48 hr) and after 72 hr, the survival rate has decreased in both cell lines and its reduction trend is significant ($P < 0.0001$) (Fig. 2, 72 hr).




Fig. 1. Growth inhibitory effect of mitomycin C on L929 and TCC cells. Cells were grown in 96-well plates and incubated for 24, 48, or 72 hr with 0, 25, 50, 100, 200, or 400 µg/mL of mitomycin C. Cell viability was measured with MTT assays. Mean $\pm$ SD, ANOVA, $n=3$. ($P<0.05^*$, $P<0.01^{**}$, $P<0.001^{***}$, $P<0.0001^{****}$).

MTT results 24, 48, 72 hr after adding mannan to L929 and TCC cells

After 24 hr, viability was significantly less in the TCC than in the L929 cells at mannan concentrations of 623, 1250, 2500, and 5,000 µg/mL (Fig. 3, 24 hr). The greatest decrease in TCC cell viability was at the lowest mannan concentration.

After 48 hr, viability was significantly less in the TCC than in the L929 cells only at 5,000 µg/mL mannan. No significant differences were seen at any other mannan concentrations (Fig. 3, 48 hr). After 72 hr no viability differences were seen between the two cell lines at any mannan concentrations (Fig. 3, 72 hr).




Fig. 2. Growth inhibitory effect of mannan on L929 and TCC cells. Cells were grown in 96-well plates and incubated for 24, 48, or 72 hr with 0, 625, 1250, 2500, 5000 or 10000 µg/mL of mannan. Cell viability was measured with MTT assays. Mean $\pm$ SD, ANOVA, $n=3$. ($P<0.05^*$, $P<0.01^{**}$, $P<0.0001^{****}$).

Flow cytometry

Apoptosis was analyzed after 72 hr by flow cytometry. Table 2 shows the effects of mannan, mitomycin C, and conjugate samples at the concentrations listed. In this data, live cells that were negative for both Annexin-V and PI appear in Q4. Cells in Q3 were in early apoptosis and Annexin-V-positive and PI-negative. Cells in Q2 were in late apoptosis; their cell walls are slightly permeable, and they are positive for both Annexin V and PI. Cells in Q1 are necrotic or dead and stained only


with PI. Apoptosis, shown in Q3+Q2, was induced by mannan in both cell lines at similar levels of 41.5 and 41% for L929 and TCC, respectively. Mitomycin C at 25 µg/mL induced apoptosis in 62.1% of L929 cells and 43.5% of TCC cells. Apoptosis was greater in cells treated with 400 µg/mL of mitomycin C than in those treated with 25 µg/mL. Apoptosis was less in the L929 conjugate-treated cells than in those treated with mannan or mitomycin C and similar in the TCC cells.

Table 1. The effects of the conjugate, mitomycin C, and mannan samples on L929 and TCC cells after 24,48, and 72 hr

Conjugate													
Cell Line		L929						TCC					
Concentration (µg/ml)	0	43.75	87.5	175	350	700	0	43.75	87.5	175	350	700	
Time post incubation	24 hr	100±0	92088±14775	100249±10992	100661±13214	87227±7212	8579±6508	100±0	101171±8335	111046±16419	38107±12122	102874±1332	103529±5596
	48 hr	100±0	112019±6570	106269±11057	113778±1980	96011±2485	62127±7408	100±0	83183±1001	83227±190	73049±0.791	52198±0.299	32813±4033
	72 hr	100±0	72831±6916	69074±5373	62997±2941	56132±4625	45550±3284	100±0	79614±6738	6717±3804	50189±4830	38367±3564	21404±5123
Mitomycin C													
Cell Line		L929						TCC					
Concentration (µg/ml)	0	25	50	100	200	400	0	25	50	100	200	400	
Time post incubation	24 hr	100±0	67855±2468	59483±4249	43784±1972	11478±0.182	10955±0.632	100±0	64455±9.148	58444±6.188	44372±2.715	37841±4.833	24975±3260
	48 hr	100±0	18582±4369	11426±1.208	7475±0.597	7194±0.753	7383±0.466	100±0	25933±4.066	20630±1.657	12688±1.451	12277±1.139	12381±1.181
	72 hr	100±0	7324±0.329	6180±0.235	6281±0.106	98202±7.219	6571±0.201	100±0	18667±2.940	12824±2.389	10200±1.633	9942±1.518	10351±1.351
Mannan													
Cell Line		L929						TCC					
Concentration (µg/ml)	0	625	1250	2500	5000	10000	0	625	1250	2500	5000	10000	
Time post incubation	24 hr	100±0	103060±7.487	102716±2.143	100159±10.166	104101±0.650	104955±5.459	100±0	75091±3.009	84913±7.461	78490±0.970	88319±7.874	10857±2.338
	48 hr	100±0	88854±14.789	94003±7.880	117088±16.798	129742±4.898	114220±14.312	100±0	98699±5.549	107982±1.635	109247±1.306	106361±1.818	101062±3.313
	72 hr	100±0	101900±25.35	106785±14.380	100643±2.357	98202±7.219	106197±2.989	100±0	98913±9.736	91708±13.525	104441±13.309	104048±17.419	10198±5.39

Table 2. Flow cytometry analysis of apoptosis induced by the mannan, mitomycin C, and conjugate samples on TCC and L929 and TCC cells after 72 hr

		Untreated (%)	Mannan 1250 µg/ml (%)	Mitomycin C 25 µg/ml (%)	Mitomycin C 400 µg/ml (%)	Conjugate 700 µg/ml (%)
72 hr' treatment	L929 Cell Line	Vital population (Q4)	72.5	53.5	34.5	68.5
		Early apoptosis (Q3)	14.4	16.8	50.2	13.0
		Late apoptosis (Q2)	11.4	24.7	11.9	13.6
		Necrosis (Q1)	1.75	4.94	3.45	4.84
72 hr' treatment	TCC Cell Line	Vital population (Q4)	71.1	51.1	42.9	57.4
		Early apoptosis (Q3)	13.6	16.5	21.7	27.0
		Late apoptosis (Q2)	13.4	24.5	39.6	15.6
		Necrosis (Q1)	1.88	7.91	2.39	0.04

**Fig. 3.** Flow cytometry results of L929 and TCC cells following treatment with mannan, mitomycin C at 25 and 400 µg/ml, and mannan-mitomycin C conjugate. Untreated cells served as control.

Morphological alterations

After 72 hr the morphology of L929 and TCC cells treated with the highest concentration of each compound was observed with inverse optical microscopy L929 cells treated with the conjugate

or mitomycin C had fewer cells than controls and cell shrinking, while those treated with mannan showed a population increase and no cell degradation (Fig. 5). After 72 hr conjugate-treated and mitomycin C-treated TCC cells showed

decreased cell numbers, cytoplasmic granulation, and rounding, while those treated with mannan showed no substantial morphological changes (Fig. 6).



Fig. 5. Microscopy images of L929 cells treated with mannan-mitomycin C conjugate, mitomycin C, and mannan. L929 cells were incubated for 72 hr with 700 $\mu\text{g/mL}$ mannan-mitomycin C conjugate, 400 $\mu\text{g/mL}$ mitomycin C, and 10,000 $\mu\text{g/mL}$ mannan. (a) untreated cells; (b) conjugate-treated cells; (c) mitomycin C-treated cells; (d) mannan-treated cells (10X).



Fig. 4. Microscopy images of TCC cells treated with mannan-mitomycin C conjugate, mitomycin C, and mannan. TCC cells were incubated for 72 hr with 700 $\mu\text{g/mL}$ mannan-mitomycin C conjugate, 400 $\mu\text{g/mL}$ mitomycin C, or 10,000 $\mu\text{g/mL}$ mannan. (a) untreated cells; (b) conjugate-treated cells; (c) mitomycin C-treated cells; (d) mannan-treated cells (10X).

Discussion

To date, several methods have been used to treat various cancers, often with undesirable side effects and low response to treatment. Hence, researchers continue to search for new and more effective ways to control cancer with fewer side effects. Numerous studies have analyzed the effect of yeast extract on different types of cancers (19, 20). Research on fungal and yeast polysaccharides has shown their

significant effects in terms of immunopotentiality and the ability to control carcinogenesis (18, 21). One polysaccharide with this feature is mannan in the cell wall of *S. cerevisiae*, a yeast known to be non-toxic in humans (22). This polysaccharide stimulates the immune system to produce cytokines that bind TLR-4 receptors on the surface of monocytes. Mitomycin C, an anti-cancer antibiotic, is activated and converted into an alkylating agent through an enzyme-mediated reduction, which causes DNA cross-linking (17). Mitomycin C is widely used to treat bladder cancer, which is why we chose it in this study. Tolley et al. reported that mitomycin C treatment reduced the number of subsequent returns and increased recurrence intervals in bladder cancer patients (23). Numerous studies on the effects of mitomycin C have been conducted including those by Bouffieux et al. (1995), Solsona (1999), Savino (2010), and James (2012), illustrating its importance in bladder cancer treatment (24-27). In this study the TCC cells were used because they express TLR-4 and thus can be specific targets for mannan (11, 12). Several cancer cell types express TLR-4, including those of the colon, stomach, prostate, breast, ovary, and brain (14, 15). The aim of this study was to investigate the inhibitory and apoptotic effects and selectivity of a mannan-mitomycin C conjugate, and these two compounds individually on L929 and TCC cells.

Inhibitory effect of mannan-mitomycin C conjugate, mitomycin C and mannan on L929 and TCC cell lines

The mannan-mitomycin C conjugate demonstrated its selective and inhibitory effects on L929 and TCC cells after 48 and 72 hr of treatment (Fig. 1). After 72 hr, the selective effect of the conjugate on the two cell lines at concentrations of 175, 350, and 700 $\mu\text{g/mL}$ was statistically significant. Mitomycin C alone had a time and dose -dependent inhibitory effect on both cell lines, and the normal cells were more sensitive to mitomycin C treatment than cancer cells.

Mannan alone had little inhibitory effect on either cell type and even increased to some extent the proliferation of normal cells after 48 and 72 hr of treatment. However, its growth-inhibitory effect on cancer cells was seen after 24 hr.

Apoptotic effect of mannan-mitomycin C conjugate, mitomycin C and mannan on L929 and TCC cell lines

Flow cytometry showed that the mannan-mitomycin C conjugate induced (early and late) apoptosis in both L929 and TCC cells. This effect was about two times greater on the TCC than on the L929 cells (Table 3). Mitomycin C alone induced apoptosis in both cell types at both 25 and 400 µg/mL in a dose-dependent manner. This effect was greater in the L929 than in the TCC cells (Table 2).

It is likely that the mannan-mitomycin C conjugate binds TCC cell TLR-4 receptors via the mannan moiety(14, 15). The mitomycin C moiety of the conjugate is likely to fragment the cell's DNA (28, 29), causing the level of cell cycle proteins such as P53 and P21 to increase (29, 30), and prevent the proliferation of cancer cells. If these proteins cannot induce repair of the damaged cell, apoptosis occurs. Mitomycin C, through the internal apoptosis pathway, reduces the level of anti- apoptotic proteins, such as Bcl-2, and activates Bax and Bad pro-apoptotic proteins, which translocate to the mitochondria. This changes the mitochondrial membrane potential, releasing cytochrome c from

the mitochondria, from where it translocates to the cell cytoplasm. Cytochrome c, with ATP and dATP, causes oligomerization of Apaf-1, and procaspase 9 attaches to them to form the apoptosome. Finally, caspase 9 is activated and a caspase cascade begins, eventually leading to cell death (30, 31). Our study suggests that the mannan-mitomycin C conjugate may be more specific for bladder cancer cells than mitomycin C alone.

In this study the inhibitory and apoptotic effects of the mannan-mitomycin C conjugate was greater on cancer cells than on normal cells while mitomycin C was more cytotoxic to normal cells than cancer cells. These results may indicate conjugate selectivity for cancer cells due to the mannan moiety binding to TLR-4 on TCC cells. This was an in vitro study, but the results suggest that its effect on animals should be investigated.

Acknowledgment

We acknowledge the Immunogenetic Center at Bu-Ali Research Institute, Mashhad University of Medical Sciences for the financial support (Grant Number: 930757). The authors declare no conflict of interest.

References

1. Vikram R, Sandler CM, Ng CS. Imaging and staging of transitional cell carcinoma: part 1, lower urinary tract. *American Journal of Roentgenology*. 2009;192(6):1481-7.
2. Asgari M, Ebrahimi M, Farshidpour M. comparison of transitional cell carcinoma grading of bladder according to urine cytology and bladder biopsy. *Razi Journal of Medical Sciences*. 2003;10(35):427-31.
3. Burger M, Catto JW, Dalbagni G, Grossman HB, Herr H, Karakiewicz P, et al. Epidemiology and risk factors of urothelial bladder cancer. *European urology*. 2013;63(2):234-41.
4. Jankovic S, Radosavljevic V. Risk factors for bladder cancer. *Tumori*. 2007;93(1):4.
5. Prout Jr GR, Barton BA, Griffin PP, Friedell GH. Treated history of noninvasive grade 1 transitional

- cell carcinoma. The National Bladder Cancer Group. *The Journal of urology*. 1992;148(5):1413-9.
6. Yazdani M, Rafati A, Zargham M, Sehri S, Khorami M. The evaluation of tumor recurrence markers P53 and CD44 in low risk superficial transitional cell carcinoma of the bladder. 2009.
7. Alexandroff AB, Jackson AM, O'donnell MA, James K. BCG immunotherapy of bladder cancer: 20 years on. *The Lancet*. 1999;353(9165):1689-94.
8. Zlotta AR, Van Vooren JP, Denis O, Drowart A, Daffé M, Lefèvre P, et al. What are the immunologically active components of Bacille Calmette- Guérin in therapy of superficial bladder cancer? *International journal of cancer*. 2000;87(6):844-52.
9. Stewart GG. *An Introduction to Brewing Science & Technology: Series III: Brewer's Yeast*. Institute of Brewing; 1998.

10. Aslani N, Nabili M, Zahedi N, Vaezi A, Badali H. The role of pattern recognition receptors (PRR) in Human Candidiasis. *Journal of Mazandaran University of Medical Sciences*. 2014;24(116):224-40.
11. Tada H, Nemoto E, Shimauchi H, Watanabe T, Mikami T, Matsumoto T, et al. *Saccharomyces cerevisiae*- and *Candida albicans*- derived mannan induced production of tumor necrosis factor alpha by human monocytes in a CD14- and Toll- like receptor 4- dependent manner. *Microbiology and immunology*. 2002;46(7):503-12.
12. Olbert PJ, Kesch C, Henrici M, Subtil FS, Honacker A, Hegele A, et al., editors. TLR4-and TLR9-dependent effects on cytokines, cell viability, and invasion in human bladder cancer cells. *Urologic Oncology: Seminars and Original Investigations*; 2015: Elsevier.
13. Bäckhed F, Meijer L, Normark S, Richter- Dahlfors A. TLR4- dependent recognition of lipopolysaccharide by epithelial cells requires sCD14. *Cellular microbiology*. 2002;4(8):493-501.
14. Andreani V, Gatti G, Simonella L, Rivero V, Maccioni M. Activation of Toll-like receptor 4 on tumor cells in vitro inhibits subsequent tumor growth in vivo. *Cancer research*. 2007;67(21):10519-27.
15. Huang B, Zhao J, Li H, He K-L, Chen Y, Mayer L, et al. Toll-like receptors on tumor cells facilitate evasion of immune surveillance. *Cancer research*. 2005;65(12):5009-14.
16. Snodgrass RG, Collier AC, Coon AE, Pritsos CA. Mitomycin C Inhibits Ribosomal RNA a Novel Cytotoxic Mechanism for Bioreductive Drugs. *Journal of Biological Chemistry*. 2010;285(25):19068-75.
17. Katzung BG, Masters SB, Trevor AJ. *Basic & clinical pharmacology*: McGraw-Hill Medical; 2016.
18. Aizawa K, Matsumoto T, Tsukada K, Ito A, Sato H, Suzuki S, et al. Antitumor effect of a baker's yeast mannan—mitomycin C conjugate against mouse hepatoma, MH134, in vivo and in vitro. *International journal of immunopharmacology*. 1989;11(2):191-5.
19. Rajan T, Benlurvankar V, Vincent S. *Saccharomyces Cerevisiae*-Induced Apoptosis of Monolayer Cervical Cancer Cells. *Asian J Pharm Clin Res*. 2017;10(8):63-6.
20. Bonyadi F, Nejati V, Tukmechi A, Hasanzadeh S, Riki M. Evaluation of Apoptotic Effects of Cell Wall and Cytoplasmic Extract from *Saccharomyces cerevisiae* on K562 Cell Line. *Armaghane danesh*. 2013;18(9):699-710.
21. Okawa Y, Ozeki Y, Suzuki K, Sakai K, Suzuki S, Suzuki M. Correlation between tumor cell cytotoxicity and serine protease activity of peritoneal macrophages from mice treated with bakers' yeast mannans. *Chemical and pharmaceutical bulletin*. 1987;35(3):1138-43.
22. Ghoneum M, Wang L, AGRAWAL S, Gollapudi S. Yeast therapy for the treatment of breast cancer: a nude mice model study. *In vivo*. 2007;21(2):251-8.
23. Tolley D, Pamar M, Grigor K-M, Lallemand G. The effect of intravesical mitomycin C on recurrence of newly diagnosed superficial bladder cancer: a further report with 7 years of followup. *The Journal of urology*. 1996;155(4):1233-8.
24. Di Stasi SM, Valenti M, Verri C, Liberati E, Giurioli A, Leprini G, et al. Electromotive instillation of mitomycin immediately before transurethral resection for patients with primary urothelial non-muscle invasive bladder cancer: a randomised controlled trial. *The lancet oncology*. 2011;12(9):871-9.
25. James ND, Hussain SA, Hall E, Jenkins P, Tremlett J, Rawlings C, et al. Radiotherapy with or without chemotherapy in muscle-invasive bladder cancer. *New England Journal of Medicine*. 2012;366(16):1477-88.
26. Solsona E, Iborra I, Ricos J, Monros J, Casanova J, Dumont R. Effectiveness of a single immediate mitomycin C instillation in patients with low risk superficial bladder cancer: short and long-term followup. *The Journal of urology*. 1999;161(4):1120-3.

27. Bouffieux C, Kurth K, Bono A, Oosterlinck W, Kruger CB, De Pauw M, et al. bladder cancer: intravesical adjuvant chemotherapy for superficial transitional cell bladder carcinoma: results of 2 European organization for research and treatment of cancer randomized trials with mitomycin C and doxorubicin comparing early versus delayed instillations and short-term versus long-term treatment. *The Journal of urology*. 1995;153(3):934-41.
28. Sasaki M, Okamura M, Ideo A, Shimada J, Suzuki F, Ishihara M, et al. Re-evaluation of tumor-specific cytotoxicity of mitomycin C, bleomycin and peplomycin. *Anticancer research*. 2006;26(5A):3373-80.
29. Kang SG, Chung H, Yoo YD, Lee JG, Choi YI, Yu YS. Mechanism of growth inhibitory effect of Mitomycin-C on cultured human retinal pigment epithelial cells: apoptosis and cell cycle arrest. *Current eye research*. 2001;22(3):174-81.
30. Wu K-Y, Wang H-Z, Hong S-J. Mechanism of mitomycin-induced apoptosis in cultured corneal endothelial cells. *Molecular vision*. 2008;14:1705.
31. Koff JL, Ramachandiran S, Bernal-Mizrachi L. A time to kill: targeting apoptosis in cancer. *International journal of molecular sciences*. 2015;16(2):2942-55.