

The Role of Interleukin-4 and 13 Gene **Polymorphisms in Allergic Rhinitis:**

A Case Control Study

Afshin Shirkani^{#1}, Atena Mansouri^{#2,3}, Reza Farid Hosseini⁴, Farahzad Jabbari Azad⁴, Reihaneh Alsadat Mahmoudian⁵, Mehdi Montazer⁶, Abdolreza Samimi⁷, Amir Abbas Momtazi-Borojeni^{8,9}, Mohammad Reza Abbaszadegan*¹⁰, Mehran Gholamin*¹¹

Abstract

Background: Allergic Rhinitis (AR) is an IgE-mediated inflammatory disorder with high morbidity rates. The eitiology of this disease is understood to occur from a complex interaction between genetic and environmental factors. T helper type 2 cells have been shown to have a crucial role in atopic disease due to their production of the cytokines, intelukin (IL)-13 and IL-4, involved in inflammation. Research has shown single nucleotide polymorphisms (SNP) of the IL-13 and IL-4 genes to be associated increased levels of IgE and with allergic diseases such as, allergic rhinitis, asthma, and atopic dermatitis. Specifically, the rs2243250 SNP of IL-4 and the rs20541 SNP of *IL-13* have been shown to be associated with AR.

Methods: A case-control study was designed to investigate the relationship between the two SNPs rs2243250 and rs20541 with the incidence of AR. The SNPs were examined in patients with AR and healthy controls (86 patients and 86 controls). Blood samples were collected and DNA was extracted to evaluate the SNPs by RFLP-PCR.

Results: Recessive analysis model of the IL-13 gene (GG vs. AA+AG) revealed that the GG genotype was more common in AR patients (P=0.36) (OR=0.8 [81% CI 0.38-1.6]). For the IL-4 gene (TC vs. TT+CC), the TC genotype was more common in AR patients (P = 0.0022)(OR = 0.71 [60% CI 1.41-5.02]). Furthermore, in the IL-4 gene, the 590 T>C polymorphism had a significant association with AR. However, no association was found between AR and the IL-13 rs20541 polymorphism.

Conclusions: Our findings suggest that the IL-13 polymorphism (rs20541, Exo 4, G>A, Arg130Gln) and IL-4 polymorphism (rs2243250= C-590T, promoter, T>C) are co-associated with AR and sensitivity to aeroallergens. However, this study used a cohort of AR patients and healthy controls from the northeast of Iran. Given the influence of ethnicity and environment on genetics, further investigation is needed to elucidate the role of SNPs in IL-4 and IL-13 in AR among different populations.

Keywords: Allergic rhinitis, Interleukin 4, Interleukin 13, Single nucleotide polymorphism.

- 1: Allergy and Clinical immunology Department, School of Medicine, Bushehr University of Medical Science, Bushehr. Iran.
- 2: Nanotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran.
- 3: Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran.
- 4: Allergy Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.
- 5: Immunology Research Center, Mashhad University of medical Sciences, Mashhad, Iran.
- 6: Department of Pathology Shiraz University of Medical Sciences, Shiraz, Iran.
- 7: Department of Anesthesiologie and prehospital emergency and intensivmedicine and pain, St. Et. Elisabeth Hospital Dorsten, KKRN, Westfalen Wilhelm University Munster, Germany.
- 8: Student Research Committee, Nanotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.
- 9: Department of Medical Biotechnology, Student Research Committee, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.
- 10: Medical Genetics Research Center, Faculty of Medical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran
- 11: Department of Laboratory Sciences, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran. #The first and the second authors contributed equally to this work.
- *Corresponding authors: Mehran Gholamin; Tel: +98 51 37112343, Fax: +98 51 37112343; E-mail: GholaminM@mums.ac.ir & Mohammad Reza Abbaszadegan; Tel: +98 51 37112343, Fax: +98 51 37112343; E-mail: Abbaszadeganmr@mums.ac.ir. Received: Sep 23, 2018; Accepted: Dec 27, 2018

Introduction

Allergic rhinitis (AR) is a highly prevalent inflammatory disorder impacting the upper respiratory tract (1, 2). The development of AR occurs from a complex interaction between environmental and genetic factors. The presence of AR is characterized by elevated levels of plasma immunoglobulin (Ig)E and an overexpression of T helper type 2 (Th2) cytokines (3).

Although the etiology remains to be clearly elucidated, the pathogenesis of AR has been found to involve different inflammatory cells including, mast cells, CD4+T cells, B cells, macrophages, and eosinophils infiltrating the nasal cavity following exposure to a stimulating allergen (4). In AR, dysregulation of the Th1/Th2 immune response leads to the production and binding of allergenspecific immunoglobulin IgE to mast cells (5, 6). When the upper airway is exposed to allergens, the allergens interact with the IgE antibodies which leads to the release of a number of cytokines, chemokines and inflammatory mediators by mast cells. This leads to the accumulation of T cells, basophils, mast cells, and eosinophils within the nasal cavity (5-7). The release of cytokines, specifically, interleukin (IL)-4 and IL-13, are known to hold a key role in the pathogenesis of AR (4, 8).

Genetic factors have been found to have a strong association with the development and severity of AR, and response to treatment (3, 9, 10). Chromosome 5q31.1 has been largely investigated as this region of the genome contains several candidate genes for allergic disease, including the gene for the Th2 cytokines, IL-4 and IL-13 (11). allergic rhinitis, Within the Th2 response predominantly entails T cells infiltrating the nasal cavity and releasing IL-4 and IL-13. The release of IL-4 and IL-13 significantly contributes to the IgEmediated allergic inflammation (5). IL-4 is a B cell growth factor known to have a large role in the allergic response. Here, IL-4 and IL-13 is involved in promoting isotype class switching from IgM to IgE, which encourages T cell differentiation towards a Th2 cell phenotype, and enhances the antigen presenting capacity of B cells (12-14). In addition, IL-13 shares many biological activities with IL-4 and holds a critical role in the allergic response as it is also involved in isotype class switching to IgE.

However, the release of *IL-13* by Th2 lymphocytes has a role in modulating the inflammatory allergic response and promotes tissue remodeling in an allergic diseases similar to AR (15).

Single nucleotide polymorphisms (SNPs) in both the IL-4 and IL-13 genes have been reported to be associated with allergic responses (15-21). With respect to IL-4, the C-590T (rs2243250) SNP located in the promoter region has been previously shown to correlate with an increased risk of AR. This SNP is understood to up-regulate IL-4 gene expression and subsequently increase the levels of plasma IgE, exacerbating the symptoms of AR (17-20). Furthermore, the IL-13 gene has several SNPs which are often associated with allergic disorders (21-24). Seven polymorphisms have been reported within the IL-13 gene however the SNP, Arg130Gln (rs20541), located in exon 4 of the IL-13 gene, has been identified to have the strongest association with high levels of plasma IgE (21,24) and with the development of AR (19, 21, 25).

In this study, we assumed that the functional polymorphisms in *IL-4* and *IL-13*, rs2243250 and rs20541, respectively contribute to the susceptibility of AR. To test this, we performed a genotype analysis for the C-590T SNP in *IL-4* and the Arg130 Gln SNP in *IL-13* in a cohort of patients from the north east region of Iran.

Materials and methods

Study design and subjects

This study was approved by the Ethical Committee of Mashhad University of Medical Sciences. The case-control study included 86 patients with allergic rhinitis and 86 healthy subjects. AR was confirmed by the skin prick test, and diagnosed according to the Allergic Rhinitis and Impact on Asthma (ARIA) guideline criteria for all participants (2). A questionnaire was provided to each participant, including signs and symptoms of allergic rhinitis based on the ARIA guidelines. Both groups were asked to read and sign informed consent forms. Following this, clinical examination and skin-prick tests were conducted to determine the presence of allergies. Specifically, the skin-prick test, contained extracts of common allergens including aeroallergens (such as pollen, trees, grasses and weeds) and dust mites, was performed to ensure the diagnosis of allergy in all the participants. The skin prick tests were negative in the control group. The control group consisted of relatives of patients, staff in the **Immunology** department, students, volunteers. Healthy subjects were placed in the control group. Individuals with possible chronic systemic disorder, atopy and allergic disease, a family history of asthma, allergy and airway diseases were excluded from the control group. The principles of the Helsinki Convention apply to this study and no additional cost was required from the patients. The inclusion criteria were as follows for the AR patients: a positive history of allergic rhinitis, a positive physical examination, and positive skin-prick test. Patients with a diagnosis of allergic rhinitis, who referred to the allergy clinic of Ghaem Hospital, were included in the study. Patients with malignant disease, auto-immune disease, lung disease, psoriasis or

any diffuse dermatitis or respiratory infection 30 days prior to recruitment for the study were excluded from the patient group.

DNA extraction

After confirming the presence or absence of allergic retinitis in the participants, 5ml of venous blood was collected from both the patient and control groups in falcon tubes containing EDTA. DNA was extracted using an extraction kit (QIAGEN Germany Cat No. /ID: 69504).

Genotyping

Genotyping of the two SNPs, rs2243250 of the *IL-4* gene and Andrs20541 of the *IL-13* gene, was performed using polymerase chain reaction (PCR) followed by restriction fragment length polymorphism (RFLP). Characteristics of the SNPs, primer sequences, PCR products size and restriction enzymes are provided in Table 1.

Table 1. The characteristics of SNPs, primer sequences, PCR product size and restriction enzymes

Name of SNP	Primer sequence	PCR product (bp)	Restriction Enzyme
IL-4 (rs2243250)	(F)= 5'-CTTCCGTGAGGACTGAATGAGAC-3' ®=5'GCAAATAATGATGCTTTCGAAGTTTCAG3'	195	AvaII
IL-13 (rs20541)	(F)=5'-TAAACTTGGGAGAACATGGT-3' (R)=5'-TGGGGAAAGATAGAGTAATA-3'	236	NLAIV

PCR was carried out in a thermal cycler in a total volume of 20 µl. This mixture contained 1 µl of each primer, 1 µl of the extracted DNA, 2 µl MgCl₂, 0.4 µl of each deoxynucleotide triphosphate, 0.25 µl of Taq DNA polymerase, 2 μl of 10X buffer and 12.35 μl of nuclease-free water. The genotype of the rs2243250 polymorphism was from type CC, CT, and TT, genotype of while the the rs20541 polymorphism was from type AA, CC and AC. The DNA was denatured at 93 °C for 3 min, and temperature cycling was set at 93 °C for 30s, 57 °C (P1) or 62 °C (P2) for 30s, and 72 °C for 2 min, followed by a final extension at 72 °C for 10 min. The sizes of the generated PCR products were 100 bp. The PCR products were cleaved by AvaII for rs2243250 and NLAIV for rs20541 as follows: 25 µl of the PCR product was added to 0.5 µl of the enzyme, and then 2.5 µl of buffer

and 5 µl of PCR product was added to 17 µl of sterile distilled water, then incubated at 37 °C for 1 hour prior to running *Polyacrylamide gel* electrophoresis. The digestion produced 210 bp and 26 bp fragments for rs20541, and produced 175 bp and 20 bp for rs2243250.

Statistical analysis

The genotype and allele frequencies were calculated through chi-square test and data analysis was carried out using SPSS software (version 16). The proportion of *IL-4* and *IL-13* polymorphisms among the patients and controls was compared using the exact version of McNemar's test for matched data. Odds ratios (comparing the odds of IL-4 and IL-13 negatives among cases versus controls) and associated 95% confidence intervals were computed through an exact logistic regression.

Results

The average age of patients in this study was 26.6 years (ranging from 15 to 60 years old), the average age in the control group was 27.66 (ranging from 15 to 60 years old). In the patient group, 36 (42%) were males, while 40 (47%) individuals in the control group were males. There was no significant difference between the two groups regarding age or gender (P > 0.05). All patients in this study had a history of AR. The presence or absence of allergy was determined by

the skin-prick test in all participants. For all individuals in the control group, the test result was negative and the test was determined positive for all subjects in the patient group. There were 44 patients who were sensitive to tree pollen (51%) and 75 patients (79%) were also allergic to grass. Additionally, 82 (95%) participants were allergic to weeds, and 40 were allergic to dust (45%). Patient characteristics are presented in Table 2.

Table 2. Demographic information for the subjects in the patient and control group

Variable	Case group Number 86(%)	Control group Number 86 (%)
Sex		
Male	36 (42%)	40 (43%)
Female	50 (58%)	46 (54%)
Age (year) Minimum	5	15
Maximum	65	60
Average	25	27
Family history of asthma	50 (58%)	-
Skin prick test		
Trees	44 (51%)	-
Grass	75 (79%)	-
Weeds	82 (82%)	-
mites	40 (40%)	-

Frequency of alleles and genotypes of IL-13 and genes

Enzymatic cleavage of the 195 bp PCR product of the *IL-4* promoter region resulted in two fragments of 175bp and 20 bp. The restriction site is eliminated by C allele. For exon 4 of *IL-13* gene, the 236 bp PCR product was cleaved into two fragments of 210 bp and 26 bp. The restriction site is eliminated by G allele.

The distribution of *IL-4* and *IL-13* genotypes, and allele frequencies of the *IL-4* SNP rs2243250 and *IL-13* SNP rs20541, were analyzed in the patient and control groups (Table 3). Frequency of the rs2243250 (C) allele in AR patients was found to be increased by 39.16% when compared to controls. Furthermore, the distribution of the CC genotype was significantly higher (23.6%) in AR patients compared to controls.

Table 3. Allele and genotypes frequencies of the IL-4 and IL-13 genes in patient with allergic rhinitis and controls.

SNP	Allele Genotype	Case	Control	PV	Odds ratio (95%CI)
rs2243250 T>C	C T CC TC TT	47 (28%) 125 (72%) 8 (10%) 31 (36%) 47 (54%)	(69%) 52 (31%) 34 (40%) 52 (60%) 0 (0)	0.0001 0.001 0.0001 0.0022 0.0001	6.3 (3.8-9.7) 0.1 (0.1-0.2) 6.3 (2.7-11.8) 0.71 (1.41-5.02) 0
rs20541 G>A	G A GG AG	150 (86%) 22 (14%) 70 (81%) 16 (19%)	(80%) 27 (20%) 66 (76%) 20 (24%)	0.23 0.24 0.36 0.28	0.76 (0.42-1.39) 1.31 (0.71-2.2) 0.8 (0.38-1.66) 1.32 (0.63-2.7)

The C allele in rs2243250 was associated with an increased the risk of AR by 6.3-fold (95%CI: 3.8-9.7). The CC genotype of rs2243250 in *IL-4* was associated with a 6.3-fold (95% CI: 2.7-11.8) increased risk of AR occurrence. However, the SNP rs20541 in exon 4 of the *IL-13* gene was not found to be associated with the risk of AR.

Discussion

AR is a multifaceted disorder that originates from a complex interaction of genetic and environmental factors that can lead to the dysregulation of the immune system (3, 26). The present study was aimed at evaluating the association of SNPs in the *IL-4* promoter (rs2243250; C-590T) and exon 4 of the *IL-13* gene (rs20541; Arg130Gln) with the incidence of AR in a cohort of patients from northeastern Iran.

It has been well-documented that IL-4 has a critical role in the pathogenesis of AR, particularly in the late phase of the disease (5). Mast cells and T helper cells are the main source of IL-4 promoting the differentiation of Th2 cells and IgE production (5, 27). Our results showed an association between the C-590T SNP in the IL-4 promoter gene and AR risk. However, the Arg130Gln SNP in IL-13 showed no significant association with the risk of AR. We also found that the CC genotype and C allele of the C-590T polymorphism had a positive association with the incidence of AR. These finding are in accordance with work from previous studies evaluating the relationship between IL-4 SNPs and the risk of various allergic-related diseases (18). A large meta-analysis evaluating several published human studies revealed that the T allele and TT genotype of the C-590T polymorphism in the IL-4 gene promoter have a positive association with AR risk (18).

The promoter region of the *IL-4* gene contains a 590C/T SNP that is understood to interact with nuclear transcription factors and regulate *IL-4* expression. Specifically, the T allele has been shown to enhance the binding of nuclear transcription factors to the promoter region, ultimately upregulating the *IL-4* expression (19, 28, 29). Furthermore, it was found that there is a relationship between *IL-4* -590C/T polymorphisms and IgE levels. The substitution of C with T in the -590 position of the *IL-4* promoter has been observed to

increase IgE levels in patients with asthma (30, 31). In China, AR patients with the TT polymorphism have been found to have significantly higher levels of IgE and an increased risk of AR than those with the CT/CC genotypes (19). Contrary to previous findings, Movahedi et al. reported that in AR patients from an Iranian population in Tehran, the CC genotype in the -590C/T SNP within IL-4 gene was associated with increased risk of AR. Patients with the TT genotype were found to have a negative association with AR (20). The inconsistencies may be explained by the variations in allele frequencies among different ethnic groups. Therefore, the TT genotype in the population in the north east of Iran may be more frequent than in those from Tehran. The contradictory results from our study and the Tehran study are supported by previous research that has shown similar conflicting findings. A separate study observed a protective effect for the CC genotype when compared to the TC and TT genotypes of patients with allergic-related disorders, including AR, asthma, and atopic dermatitis (32). Similarly, additional studies of patients with rhino conjunctivitis and hay fever showed no significant association between the -590C/T SNP and allergy (33, 34). Although we did not observe any association between the Arg130Gln SNP in IL-13 and a risk of AR, it has been frequently reported that this SNP is associated with plasma levels of IgE and AR risk (19, 22, 31, 35-38). Lack of an association in our study may be due to the differences in ethnic backgrounds and the effect of the environment on the genetic background of our examined population. This is supported by research from a meta-analysis showing that the contradictory findings regarding the association of specific SNPs with allergic rhinitis is likely a result of the genetic variation among different ethnicities and the effect of different environmental factors (38). The various results reported from different studies regarding genetic polymorphisms of cytokines in allergic rhinitis indicate different roles for particular genetic polymorphisms among various ethnicities. Diseases with complex traits such as allergic rhinitis are likely to be explained by complex genetic interactions, as well as the genetic and environmental factors. Therefore, different racial groups could express different polymorphisms within candidate genes.

Due to the differences among various studies examining the relationship between the genetic polymorphisms of cytokines, levels of cytokines production, and gene expression, additional research is necessary to further identify and evaluate the SNPs in genes involved in allergic rhinitis and the relationship with the serum level of inflammatory cytokines and IgE. Furthermore, it is suggested that this genetic relationship be in other Iranian evaluated races. Another suggestion would be to incorporate a greater number of clinical indices to provide a more accurate assessment of the relationship between the

References

- 1. Bourdin A, Gras D, Vachier I, Chanez P. Upper airway· 1: Allergic rhinitis and asthma: united disease through epithelial cells. Thorax. 2009;64(11):999-1004.
- 2. Shirkani A, Mansouri A, Abbaszadegan MR, Faridhosseini R, Azad FJ, Gholamin M. GATA3 Gene Polymorphisms Associated with Allergic Rhinitis in an Iranian Population. Reports of biochemistry & molecular biology. 2017;5(2):97.
- 3. Davila I, Mullol J, Ferrer M, Bartra J, Del Cuvillo A, Montoro J, et al. Genetic aspects of allergic rhinitis. J Investig Allergol Clin Immunol. 2009;19(Suppl 1):25-31.
- 4. Small P, Frenkiel S, Becker A, Boisvert P, Bouchard J, Carr S, et al. Rhinitis: A Practical and Comprehensive Approach to Assessment and Therapy. Journal of otolaryngology. 2007;36.
- 5. Pawankar R, Mori S, Ozu C, Kimura S. Overview on the pathomechanisms of allergic rhinitis. Asia Pacific Allergy. 2011;1(3):157-67.
- 6. Howarth P. Allergic rhinitis: not purely a histamine-related disease. Allergy. 2000;55(s64):7-16.
- 7. Kay A. Allergy and allergic diseases. New England Journal of Medicine. 2001;344(1):30-7.
- 8. Riccio A, Tosca M, Cosentino C, Pallestrini E, Ameli F, Canonica G, et al. Cytokine pattern in allergic and non-allergic chronic rhinosinusitis in asthmatic children. Clinical & Experimental Allergy. 2002;32(3):422-6.
- 9. Toda M, Ono SJ. Genomics and proteomics of allergic disease. Immunology. 2002;106(1):1-10. 10. Peden DB. Influences on the development of allergy and asthma. Toxicology. 2002;181:323-8.

genetic polymorphism of cytokines, and the development and severity of allergic rhinitis. These efforts will help to create improved diagnostic and medical treatments for allergic rhinitis.

Acknowledgment

This study was supported by a grant from Mashhad University of Medical Sciences (#90085). The results described in this paper were part of the fellow dissertation in Allergy and clinical Immunology.

The authors declare no conflicts of interest for this report.

- 11. Mou Z, Shi J, Tan Y, Xu R, Zhao Z, Xu G, et al. Association between TIM-1 gene polymorphisms and allergic rhinitis in a Han Chinese population. J Investig Allergol Clin Immunol. 2010;20(1):3-8.
- 12. Coffman RL, Ohara J, Bond MW, Carty J, Zlotnik A, Paul W. B cell stimulatory factor-1 enhances the IgE response of lipopolysaccharide-activated B cells. The Journal of Immunology. 1986;136(12):4538-41.
- 13. Bacharier LB, Geha RS. Molecular mechanisms of IgE regulation. Journal of Allergy and Clinical Immunology. 2000;105(2):S547-S58.
- 14. Seder RA, Paul WE. Acquisition of lymphokine-producing phenotype by CD4+ T cells. Annual review of immunology. 1994;12(1):635-73.
- 15. de Vries JE. The role of IL-13 and its receptor in allergy and inflammatory responses. Journal of Allergy and Clinical Immunology. 1998;102(2):165-9.
- 16. Yokouchi Y, Nukaga Y, Shibasaki M, Noguchi E, Kimura K, Ito S, et al. Significant evidence for linkage of mite-sensitive childhood asthma to chromosome 5q31–q33 near the interleukin 12 B locus by a genome-wide search in Japanese families. Genomics. 2000;66(2):152-60.
- 17. Yokouchi Y, Shibasaki M, Noguchi E, Nakayama J, Ohtsuki T, Kamioka M, et al. A genome-wide linkage analysis of orchard grass-sensitive childhood seasonal allergic rhinitis in Japanese families. Genes and immunity. 2002;3(1):9-13.

- 18. Li Z-p, Yin L-l, Wang H, Liu L-s. Association between promoter polymorphisms of interleukin-4 gene and allergic rhinitis risk: a meta-analysis. Journal of Huazhong University of Science and Technology [Medical Sciences]. 2014;34:306-13. 19. Lu M-P, Chen R-X, Wang M-L, Zhu X-J, Zhu L-P, Yin M, et al. Association study on IL-4, IL-13 and IL-4RA polymorphisms in mite-sensitized persistent allergic rhinitis in a Chinese population. PloS one. 2011;6(11):e27363.
- 20. Movahedi M, Amirzargar AA, Nasiri R, Hirbod-Mobarakeh A, Farhadi E, Tavakol M, et al. Gene polymorphisms of Interleukin-4 in allergic rhinitis and its association with clinical phenotypes. American journal of otolaryngology. 2013;34(6):676-81.
- 21. Wang M, Liu J, Tian X, Zhu X, Liu Y. Association of IL-13 rs20541 polymorphism and risk of allergic rhinitis: evidence from a meta-analysis. INTERNATIONAL JOURNAL OF CLINICAL AND EXPERIMENTAL MEDICINE. 2016;9(8):15914-20.
- 22. Graves PE, Kabesch M, Halonen M, Holberg CJ, Baldini M, Fritzsch C, et al. A cluster of seven tightly linked polymorphisms in the IL-13 gene is associated with total serum IgE levels in three populations of white children. Journal of Allergy and Clinical Immunology. 2000;105(3):506-13.
- 23.DeMeo D, Silverman E, Senter J, Drazen J, Barth M, Laird N, et al. Univariate and multivariate family-based association analysis of the IL-13 ARG130GLN polymorphism in the Childhood Asthma Management program. Genetic epidemiology. 2002;23(4):335-48.
- 24. Tsunemi Y, Saeki H, Nakamura K, Sekiya T, Hirai K, Kakinuma T, et al. Interleukin-13 gene polymorphism G4257A is associated with atopic dermatitis in Japanese patients. Journal of dermatological science. 2002;30(2):100-7.
- 25. Shazia M, Kanza M, Mehwish I, Irum S, Farida A, Asifa A. IL-13 gene polymorphisms and their association with atopic asthma and rhinitis in Pakistani patients. Iranian Journal of Allergy, Asthma and Immunology. 2013;12(4):391.
- 26. Bashi DS, Dowom SA, Bazzaz BSF, Khanzadeh F, Soheili V, Mohammadpour A. Evaluation, prediction and optimization the ultrasound-assisted extraction method using response surface methodology: antioxidant and

- biological properties of Stachys parviflora L. Iranian journal of basic medical sciences. 2016;19(5):529.
- 27. Okano M, Satoskar A, Abe M, Harn D, Nishizaki K, Takeda Y, et al. Interleukin-4-independent production of Th2 cytokines by nasal lymphocytes and nasal eosinophilia in murine allergic rhinitis. Allergy. 2000;55(8):723-31.
- 28. Nakashima H, Miyake K, Inoue Y, Shimizu S, Akahoshi M, Tanaka Y, et al. Association between IL-4 genotype and IL-4 production in the Japanese population. Genes and immunity. 2002;3(2):107-9. 29. Rosenwasser L, Klemm D, Dresback J, Inamura H, Mascali J, Klinnert M, et al. Promoter polymorphisms in the chromosome 5 gene cluster in asthma and atopy. Clinical & Experimental Allergy. 1995;25(s2):74-8.
- 30. Chanez P, Wenzel SE, Anderson GP, Anto JM, Bel EH, Boulet L-P, et al. Severe asthma in adults: what are the important questions? Journal of Allergy and Clinical Immunology. 2007;119(6):1337-48.
- 31. Kim JJ, Min JY, Lee JH. Polymorphisms in the IL-13 and IL-4 receptor alpha genes and allergic rhinitis. European archives of oto-rhinolaryngology. 2007;264(4):395-9.
- 32. de Guia RM, Ramos J. The-590C/T IL-4 single-nucleotide polymorphism as a genetic factor of atopic allergy. Int J Mol Epidemiol Genet. 2010;1(1):67-73.
- 33. Miyake Y, Tanaka K, Arakawa M. Polymorphisms in the IL-4 gene, smoking, and rhinoconjunctivitis in Japanese women: The Kyushu Okinawa Maternal and Child Health Study. Human immunology. 2012;73(10):1046-9.
 34. Imboden M, Nieters A, Bircher A, Brutsche M, Becker N, Wjst M, et al. Cytokine gene
- Becker N, Wjst M, et al. Cytokine gene polymorphisms and atopic disease in two European cohorts.(ECRHS-Basel and SAPALDIA). Clinical and Molecular Allergy. 2006;4(1):1.
- 35. Maier LM, Howson JM, Walker N, Spickett GP, Jones RW, Ring SM, et al. Association of IL-13 with total IgE: evidence against an inverse association of atopy and diabetes. Journal of allergy and clinical immunology. 2006;117(6):1306-13.
- 36. Wang M, Xing Z-M, Lu C, Ma Y-X, Yu D-L, Yan Z, et al. A common IL-13 Arg130Gln single

nucleotide polymorphism among Chinese atopy patients with allergic rhinitis. Human genetics. 2003;113(5):387-90.

37. Wang DY. Genetic predisposition for atopy and allergic rhinitis in the Singapore Chinese

population. Asia Pacific Allergy. 2011;1(3):152. 38. Ying X-J, Zhao S-W, Wang G-L, Xie J, Xu H-M, Dong P. Association of interleukin-13 SNP rs20541 with allergic rhinitis risk: a meta-analysis. Gene. 2013;521(2):222-6.